

Gauge Formulation for Higher Order Gravity

R. R. Cuzinatto^{1,*}, C. A. M. de Melo^{1,2,†}, L. G. Medeiros^{1,‡} and P. J. Pompeia^{1,3§}

¹*Instituto de Física Teórica, UNESP - São Paulo State University.*

Rua Pamplona 145, CEP 01405-900, São Paulo, SP, Brazil.

²*Universidade Vale do Rio Verde de Três Corações,*

Av. Castelo Branco, 82 - Chácara das Rosas,

P.O. Box 3050, CEP 37410-000, Três Corações, MG, Brazil

³*Comando-Geral de Tecnologia Aeroespacial,*

Instituto de Fomento e Coordenação Industrial. and

Praça Mal. Eduardo Gomes 50, CEP 12228-901, São José dos Campos, SP, Brazil.

Abstract

This work is an application of the second order gauge theory for the Lorentz group, where a description of the gravitational interaction is obtained which includes derivatives of the curvature. We analyze the form of the second field strength, $G = \partial F + fAF$, in terms of geometrical variables. All possible independent Lagrangians constructed with quadratic contractions of F and quadratic contractions of G are analyzed. The equations of motion for a particular Lagrangian, which is analogous to Podolsky's term of his Generalized Electrodynamics, are calculated. The static isotropic solution in the linear approximation was found, exhibiting the regular Newtonian behaviour at short distances as well as a meso-large distance modification.

*Electronic address: cuzinatto@gmail.com

†Electronic address: cassius.anderson@gmail.com

‡Electronic address: leogmediros@gmail.com

§Electronic address: pompeia@ift.unesp.br

I. INTRODUCTION

Nowadays there are many proposals to modify gravitation in order to solve several problems as the present day accelerated expansion of the universe [1, 2], or to accommodate corrections of quantum nature which arise from the classical effective backreaction of quantum matter in a curved background [3]. Effective action is widely used in quantum field theory as a powerful method of calculation. The Podolsky generalized electrodynamics, for instance, can be viewed as an effective description of quantum correction to the classical Maxwell Lagrangian [4].

For gravitation, usually higher orders terms are introduced by means of Lagrangian contributions quadratic in the Riemann tensor and their contractions [5]. This is inspired by 1-loop corrections in the Einstein-Hilbert action in the quantized weak field approximation, or in the equivalent Feynman construction of a spin-2 field on the flat Minkowski background [6]. Besides this, at the quantum level, the S matrix for the Einstein theory is finite at one-loop level, but diverges at the two-loop order [7], which motivates the introduction of derivative terms in the Riemann tensor for the action [8].

On the other hand, recently was proposed a second order construction of gauge theories based on Utiyama's approach [9], which gives exactly the same correction terms as in the Podolsky electrodynamics, but now arising from the principle of local gauge invariance [10]. Therefore, a connection between quantum corrections and gauge higher order terms in the action was conjectured, which was proved to be fulfilled also for the effective Alekseev-Arbuzov-Baikov Lagrangian of the infrared regime of QCD [11].

Here, we analyse the gauge formulation of the gravitational field based on the framework of the second order gauge theory. The simplest gauge group is given by the Lorentz homogeneous group in the context of a Riemannian description of the gravitational field. Since the gauge field is given in such case by the local spin connection, higher order in the gauge field involves naturally the derivative of the curvature tensor. In this sense, the actual higher order gravitational lagrangian should be constructed from invariants using the covariant derivative of the Riemann tensor instead of the usual quadratic terms in curvature.

The relationship between the algebraic gauge description and the geometrical one is settled by means of the introduction of the tetrad field, and the construction of the covariant derivatives associated with the both symmetries: the local Lorentz and the global diffeo-

morphic coordinate transformations. We use Latin indexes, $a, b, \dots$, for the internal Lorentz group and greek indexes for the tangent space of the space-time manifold.

The paper is structured as follows. In section II we review some results relating gauge invariance and gravitation. The field strengths F and G of the second order treatment are introduced in section III, where they are also written in their geometrical counterparts: the Riemann curvature tensor and its covariant derivative.

Section IV deals with the possible quadratic invariants of the type F^2 and G^2 . All the possible contractions are studied and only the independent invariants are kept. This counting is made in the same spirit as the systematic selection of the independent Riemann monomials done in [12] and [13]. In the following section, section V, these invariants are shown to satisfy the identity which restricts the theories that may be called of the gauge type.

Among all invariants, we select $L_P = \frac{1}{2}h \delta^\rho R_{\rho\chi} \delta_\mu R^{\mu\chi}$, the Podolsky-like Lagrangian, for calculating the equation of motion of the gravitational field. This higher order gravity application is done in section VI. For this Lagrangian, we calculate the static isotropic solution in the linear regime at section VIB, finding the regular Newtonian potential at short scales, but with a modified potential at intermediary scales.

Final remarks are given in section VII.

II. GAUGE INTERACTION AND COVARIANCE

In 1956 Utiyama [9] has shown how to implement a gauge description for gravitational interaction with matter fields $Q^A(x)$ transforming according to

$$\delta Q^A(x) = \frac{1}{2} \varepsilon^{ab}(x) (\Sigma_{ab})^A_B Q^B, \quad (1)$$

as an implementation of the local invariance exigency of the action under continuous proper Lorentz transformations, which are characterized by the generators Σ_{ab} satisfying the operation of a typical Lie group,

$$[\Sigma_{ab}, \Sigma_{cd}] = \frac{1}{2} f_{ab, cd}^{ef} \Sigma_{ef}, \quad (2)$$

where

$$f_{ab, cd}^{ef} = \left\{ [\eta_{bc} \delta_a^e - \eta_{ac} \delta_b^e] \delta_d^f - [\eta_{bd} \delta_a^e - \eta_{ad} \delta_b^e] \delta_c^f \right\} - e \leftrightarrow f$$

are the structure constants obeying the Jacobi identity. $\varepsilon^{ab} = -\varepsilon^{ba}$ are the parameters of the local transformation. The capital latin indexes are for the components of the matter field.

It was clearly shown the need to introduce the compensating field $\omega_\mu^{ab}(x)$ transforming as a connection,

$$\delta\omega_\mu^{ef} = \frac{1}{4}\varepsilon^{ab}(x) f_{ab, cd}^{\quad ef} \omega_\mu^{cd} + \partial_\mu \varepsilon^{ef}(x) . \quad (3)$$

To ensure the covariance under coordinate transformations it was necessary to define an space-time connection whose behaviour under infinitesimal diffeomorphisms is

$$\bar{\delta}\Gamma_{\mu\alpha}^\nu = \frac{\partial\delta x^\nu}{\partial x^\lambda} \Gamma_{\mu\alpha}^\lambda - \frac{\partial\delta x^\lambda}{\partial x^\mu} \Gamma_{\lambda\alpha}^\nu - \frac{\partial\delta x^\lambda}{\partial x^\alpha} \Gamma_{\mu\lambda}^\nu - \frac{\partial^2\delta x^\nu}{\partial x^\mu\partial x^\alpha}$$

The invariance of the theory implies that the compensating field must appear through the *gauge covariant derivative*

$$D_\mu Q^A \equiv \partial_\mu Q^A - \frac{1}{2}\omega_\mu^{ab}(\Sigma_{ab})^A_B Q^B \quad (4)$$

i.e.,

$$\delta D_\mu Q^A = \frac{1}{2}\varepsilon^{ab}(\Sigma_{ab})^A_B D_\mu Q^B , \quad (5)$$

and the space-time connection must appear through the space-time covariant derivative,

$$\delta_\mu Q^{\lambda\nu} \equiv \partial_\mu Q^{\lambda\nu} + \Gamma_{\mu\beta}^\lambda Q^{\beta\nu} + \Gamma_{\mu\alpha}^\nu Q^{\lambda\alpha} , \quad (6)$$

and the total covariant derivative:

$$\nabla_\mu Q^{i\nu} = \partial_\mu Q^{i\nu} - \omega_\mu^{ib} \eta_{bk} Q^{k\nu} + \Gamma_{\mu\alpha}^\nu Q^{i\alpha} . \quad (7)$$

This total derivative must commute with the mapping to the tangent space of the manifold,¹

$$Q^{i\mu} \equiv h_j^\mu Q^{ij} , \quad Q^{ij} = h_\nu^j Q^{i\nu} , \quad (8)$$

$$\nabla_\mu Q^{i\nu} \equiv h_j^\nu \nabla_\mu Q^{ij} , \quad (9)$$

¹ Note that the action of the total derivative on a tangent space field is defined by

$$\nabla_\mu Q^i \equiv D_\mu Q^i .$$

where we have introduced the tetrad field h :

$$h^j_\nu h^\mu_j = \delta^\mu_\nu, \quad h^i_\nu h^\nu_j = \delta^i_j, \quad g_{\mu\nu} = \eta_{ij} h^i_\nu h^j_\mu, \quad \eta_{ij} = h_i^\mu h_j^\nu g_{\mu\nu}, \quad h = \sqrt{\det h^j_\mu} = \sqrt{-g}.$$

The definition (9) implies the absolute parallelism of the tetrad:

$$\nabla_\mu h^j_\alpha \equiv 0, \quad (10)$$

which can be solved for the compensating field,

$$\omega^j_{i\mu} \equiv h_i^\alpha (\delta_\mu h^j_\alpha)$$

or for the space-time connection,

$$\Gamma^\nu_{\mu\alpha} \equiv h_j^\nu (D_\mu h^j_\alpha). \quad (11)$$

We will restrict our analysis to a symmetric space-time connection in order to approach the Riemannian description. The extension to the Riemann-Cartan case is quite natural, but would imply different types of invariants as admissible Lagrangians (see discussion below).

III. GAUGE FIELD LAGRANGIAN

The basic hypothesis we will assume is: the Lagrangian for the free gauge potential depends on the field, its first and second order derivatives and $L_0 = L_0(\omega_\mu^{ef}, \partial_\nu \omega_\mu^{ef}, \partial_\rho \partial_\nu \omega_\mu^{ef})$ obeys local invariance under (3). This enable us to use the results presented elsewhere [10] to construct a gauge formulation for higher order gravitation theories.

A. The field strengths

According to the work [10], we can reexpress

$$\delta L_0 = \frac{1}{2} \frac{\partial L_0}{\partial \omega_\mu^{ef}} \delta \omega_\mu^{ef} + \frac{1}{2} \frac{\partial L_0}{\partial (\partial_\nu \omega_\mu^{ef})} \delta \partial_\nu \omega_\mu^{ef} + \frac{1}{2} \frac{\partial L_0}{\partial (\partial_\rho \partial_\nu \omega_\mu^{ef})} \delta \partial_\rho \partial_\nu \omega_\mu^{ef} \equiv 0,$$

splitting it into a set of four hierarchical equations after substituting (3) and claiming the independence of the parameters ε^{ab} and their derivatives. Three of these functional equations are used to conclude that

$$L_0 = L_0(F, G); \quad \frac{\partial L_0}{\partial \omega_\mu^{ab}} \equiv 0, \quad (12)$$

where

$$F_{\mu\nu}^{ab} = \partial_\mu \omega_\nu^{ab} - \partial_\nu \omega_\mu^{ab} - \eta_{cd} \omega_\mu^{ac} \omega_\nu^{db} + \eta_{cd} \omega_\nu^{ac} \omega_\mu^{db} \quad (13)$$

and

$$G_{\beta\rho\sigma}^{ab} = D_\beta F_{\rho\sigma}^{ab} = \partial_\beta F_{\rho\sigma}^{ab} - \eta_{fd} \omega_\beta^{af} F_{\rho\sigma}^{db} + \eta_{fd} \omega_\beta^{af} F_{\rho\sigma}^{bd} . \quad (14)$$

The remaining hierarchical equation put in terms of the gauge fields F and G ,

$$\frac{\partial L_0}{\partial F_{\rho\sigma}^{ad}} f_{bcgh}^{ad} F_{\rho\sigma}^{gh} + \frac{\partial L_0}{\partial G_{\beta\rho\sigma}^{ad}} f_{bcgh}^{ad} G_{\beta\rho\sigma}^{gh} \equiv 0 , \quad (15)$$

imposes restrictions upon the functional form eventually chosen for L_0 . Substituting the structure constants, this condition can be explicitly written as

$$\frac{\partial L_0}{\partial F_{\rho\sigma}^{ad}} [\eta_{cg} \delta_b^a - \eta_{bg} \delta_c^a] F_{\rho\sigma}^{gd} + \frac{\partial L_0}{\partial G_{\beta\rho\sigma}^{ad}} [\eta_{cg} \delta_b^a - \eta_{bg} \delta_c^a] G_{\beta\rho\sigma}^{gd} \equiv 0 . \quad (16)$$

B. Geometrical variables

In this section we will show how to interpret all objects and condition of the previous sections in terms of a geometrical point of view. From (10) we read

$$\omega_\sigma^{eg} = \eta^{gc} h_c^\alpha (\partial_\sigma h_\alpha^e - \Gamma_{\sigma\alpha}^\nu h_\nu^e)$$

and therefore the field strength F is written as

$$F_{\beta\sigma}^{eg} = \eta^{gc} h_c^\alpha h_\gamma^e [\partial_\sigma \Gamma_{\beta\alpha}^\gamma - \partial_\beta \Gamma_{\sigma\alpha}^\gamma + \Gamma_{\beta\alpha}^\nu \Gamma_{\sigma\nu}^\gamma - \Gamma_{\sigma\alpha}^\nu \Gamma_{\beta\nu}^\gamma] ,$$

where we recognize the expression of the Riemann tensor [15],

$$R_{\sigma\beta\alpha}{}^\gamma \equiv \partial_\sigma \Gamma_{\beta\alpha}^\gamma - \partial_\beta \Gamma_{\sigma\alpha}^\gamma + \Gamma_{\beta\alpha}^\nu \Gamma_{\sigma\nu}^\gamma - \Gamma_{\sigma\alpha}^\nu \Gamma_{\beta\nu}^\gamma ,$$

i.e.,

$$F_{\beta\sigma}^{eg} = \eta^{gc} h_c^\alpha h_\gamma^e R_{\sigma\beta\alpha}{}^\gamma . \quad (17)$$

The easiest way to find the geometrical counterpart of G is to apply the geometrizing relations (6,8):

$$h_a^\mu h_b^\nu G_{\beta\rho\sigma}^{ab} = h_a^\mu h_b^\nu D_\beta F_{\rho\sigma}^{ab} = \delta_\beta F_{\rho\sigma}^{\mu\nu} ;$$

$$F_{\rho\sigma}^{\mu\nu} = h_a^\mu h_b^\nu F_{\rho\sigma}^{ab}$$

and use (17). We arrive at

$$G^{ab}_{\beta\rho\sigma} = h^a_{\mu} h^b_{\nu} g^{\nu\alpha} \delta_{\beta} R_{\sigma\rho\alpha}{}^{\mu}, \quad (18)$$

which is the most natural equation one would expect in view of the relation (14) between F and G .

By means of the geometrical descriptions (17) and (18), we are able to find

$$\begin{aligned} \frac{\partial L_0}{\partial F^{ad}_{\rho\sigma}} &= \frac{\partial L_0}{\partial R_{\lambda\beta\alpha}{}^{\gamma}} \frac{\partial R_{\lambda\beta\alpha}{}^{\gamma}}{\partial F^{ad}_{\rho\sigma}} = \frac{\partial L_0}{\partial R_{\sigma\rho\alpha}{}^{\gamma}} \eta_{bd} h^b_{\alpha} h_a{}^{\gamma}, \\ \frac{\partial L_0}{\partial G^{ad}_{\beta\rho\sigma}} &= \frac{\partial L_0^{(4)}}{\partial (\delta_{\lambda} R_{\gamma\nu\alpha}{}^{\mu})} \frac{\partial (\delta_{\lambda} R_{\gamma\nu\alpha}{}^{\mu})}{\partial G^{ad}_{\beta\rho\sigma}} = \frac{\partial L_0}{\partial (\delta_{\beta} R_{\sigma\rho\alpha}{}^{\mu})} g_{\alpha\omega} h_a{}^{\mu} h_d{}^{\omega}. \end{aligned}$$

With these derivatives, the condition (15) for the gauge Lagrangian is put in the form

$$\begin{aligned} &\frac{\partial L_0}{\partial R_{\sigma\rho\beta}{}^{\theta}} [\delta^{\theta}_{\nu} g_{\gamma\lambda} - \delta^{\theta}_{\gamma} g_{\nu\lambda}] R_{\sigma\rho\beta}{}^{\lambda} + \\ &+ \left[\frac{\partial L_0}{\partial (\delta_{\beta} R_{\sigma\rho\alpha}{}^{\gamma})} g_{\nu\lambda} - \frac{\partial L_0}{\partial (\delta_{\beta} R_{\sigma\rho\alpha}{}^{\nu})} g_{\gamma\lambda} \right] \delta_{\beta} R_{\sigma\rho\alpha}{}^{\lambda} \equiv 0. \end{aligned} \quad (19)$$

This is a fundamental restriction upon the Lagrangians tentatively proposed for the theory, and it is quite useful in order to choose a specific suitable invariant.

IV. QUADRATIC LAGRANGIAN COUNTING

Our goal here is to determine all possible independent quadratic Lagrangians constructed with the field strength tensors F and G considering their various symmetries. By quadratic Lagrangians we mean invariants of the type FF or GG , but not mixed terms like FG (obviously with the proper contraction of indices). We will also compute the linear case of the Einstein-Hilbert Lagrangian.

A. First Order Invariants

The symmetries to be considered in the construction of the invariants of the type FF are those inherited from F : skew-symmetry in each pair of indices: $F^{ab}_{\mu\nu} = -F^{ba}_{\mu\nu}$ and $F^{ab}_{\mu\nu} = -F^{ab}_{\nu\mu}$. Besides these, there is another which is unveiled by the geometrical form of F , eq. (17), namely

$$R_{\sigma\beta\alpha}{}^{\gamma} + R_{\beta\alpha\sigma}{}^{\gamma} + R_{\alpha\sigma\beta}{}^{\gamma} \equiv 0,$$

the familiar first Bianchi identity met in the context of the general relativity.

Once algebra and space-time indices can be transformed into each other by means of a tetrad, we will consider a compact representation for F :

$$F_{\mu\nu}^{ab} \rightarrow F_{\mu\nu}^{ab} h_c^\mu h_d^\nu \equiv (abcd) \quad .$$

Since the Lagrangians are all of the form F^2 with all allowed orders of contractions, it is always possible to rename dummy indices in such a way that the first F will keep its indices in alphabetic order. In the table below it follows all available permutations for the second F :

	Fix. a	Fix. b	Fix. c	Fix. d
cyclic	$(abcd)$	$(bacd)$	$(cabd)$	$(dabc)$
	$(acdb)$	$(bcda)$	$(cbda)$	$(dbca)$
	$(adbc)$	$(bdac)$	$(cdab)$	$(dcab)$
non-cycl.	$(abdc)$	$(badc)$	$(cadb)$	$(dacb)$
	$(acbd)$	$(bcad)$	$(cbad)$	$(dbac)$
	$(adcb)$	$(bdca)$	$(cdba)$	$(dcba)$

(20)

By means of a change in one pair of indices, one can see that the non-cyclic permutations are all proportional to the cyclic ones. Considering only the cyclic permutations and changing two pairs of indices, the table is reduced to:

	Fix. a	Fix. b	Fix. c	Fix. d
cyclical	$(abcd)$	—	—	—
	$(acdb)$	$(bcda)$	—	—
	$(adbc)$	$(bdac)$	$(cdab)$	—

The skew-symmetries of the first F (which has been taken in alphabetic order) leads one to restrict once more the possible contractions to the three quadratic invariants

$$\begin{aligned}
I_1^F &= (abcd) (abcd) \\
I_2^F &= (abcd) (acdb) \\
I_3^F &= (abcd) (cdab) \quad .
\end{aligned}
\tag{21}$$

We now analyze invariants constructed with one trace of F . The only non-null type of trace are those obtained by contracting one index of the first pair with one index of the

second pair, in view of the skew-symmetry of this object. All possibilities are proportional to

$$Tr F \rightarrow h_c^\nu F_{\mu\nu}^{ca} h_b^\mu \equiv (\cdot ab \cdot) \quad \text{or} \quad (\circ ab \circ) .$$

The quadratic invariants are given by,

$$\begin{aligned} I_1^{TrF} &= (\cdot ab \cdot) (\circ ab \circ) \\ I_2^{TrF} &= (\cdot ab \cdot) (\circ ba \circ) . \end{aligned} \tag{22}$$

Still, one can construct a linear invariant taking a double trace of F :

$$I^{TrTrF} = h_c^\nu F_{\mu\nu}^{ca} h_a^\mu \equiv (\cdot \circ \circ \cdot) .$$

B. Second Order Invariants

Let us introduce a similar notation to the one used in the case of F , i.e.,

$$G_{\beta\rho\sigma}^{ab} \rightarrow h_c^\beta h_d^\rho h_e^\sigma G_{\beta\rho\sigma}^{ab} \equiv [abcde] ,$$

where we identify the following symmetries:

(i) antisymmetry by permutation of indices in the first pair and last two of them,

$$[abcde] = -[bacde] = -[abced] ;$$

(ii) Bianchi identity among the last three indices,

$$[abcde] + [abdec] + [abecd] = 0 .$$

1. Invariants of GG kind

The quadratic combinations are now in a larger amount than in the F^2 case. In fact, we have five tables like (20), one to each letter labeling, since we can associate

$$[abcde] = c(abde) .$$

Using the symmetries cited above, one finds that the 5! G^2 invariants are reduce to just two kinds:

$$\begin{aligned} I_1^G &= [abcde] [abcde] \\ I_2^G &= [abcde] [debac] . \end{aligned}$$

The detailed and cumbersome calculations are made in the appendix A.

2. Invariants Involving Traces

There are three independent types of traces for G :

$$\begin{aligned} T_{abc}^{(1)} &= h_d^\beta G_{\beta\rho\sigma}^{da} h_b^\rho h_c^\sigma \equiv [\cdot a \cdot bc] \\ T_{abc}^{(2)} &= h_d^\rho G_{\beta\rho\sigma}^{da} h_b^\beta h_c^\sigma \equiv [\cdot ab \cdot c] \\ T_{abc}^{(3)} &= g^{\beta\rho} G_{\beta\rho\sigma}^{ab} h_c^\sigma \equiv [ab \cdot \cdot c] . \end{aligned} \tag{23}$$

Again using symmetries (see appendix A) we arrive at:

$Tr\mathcal{G}_3 = [\cdot ab \cdot c] [\cdot a \cdot bc]$	$Tr\mathcal{G}_{11} = [\cdot ab \cdot c] [\cdot bc \cdot a]$	(24)
$Tr\mathcal{G}_5 = [\cdot ab \cdot c] [\cdot c \cdot ab]$	$Tr\mathcal{G}_{14} = [\cdot ab \cdot c] [\cdot ab \cdot c]$	
$Tr\mathcal{G}_6 = [ab \cdot \cdot c] [\cdot a \cdot bc]$	$Tr\mathcal{G}_{17} = [ab \cdot \cdot c] [ab \cdot \cdot c]$	
$Tr\mathcal{G}_{10} = [\cdot ab \cdot c] [\cdot ba \cdot c]$	$Tr\mathcal{G}_{18} = [ab \cdot \cdot c] [ac \cdot \cdot b]$	

while for double traces we have:

$$\begin{aligned} TrTr\mathcal{G}_1 &= [\cdot \circ b \cdot \circ] [\cdot \circ b \cdot \circ] \\ TrTr\mathcal{G}_2 &= [\circ b \cdot \cdot \circ] [\circ b \cdot \cdot \circ] \\ TrTr\mathcal{G}_3 &= [\cdot \circ b \cdot \circ] [\circ b \cdot \cdot \circ] . \end{aligned} \tag{25}$$

C. Bianchi Identities

As we already said, until now we have not used the first Bianchi identity:

$$R^{\sigma\chi\rho\beta} + R^{\chi\rho\sigma\beta} + R^{\rho\sigma\chi\beta} \equiv 0 . \tag{26}$$

In geometrical variables, the cyclic property of G is translated to the second Bianchi identity:

$$\delta^\mu R^{\sigma\chi\rho\beta} + \delta^\sigma R^{\chi\mu\rho\beta} + \delta^\chi R^{\mu\sigma\rho\beta} \equiv 0 .$$

These identities reduce the number of independent invariants, since $F \propto R$ and $G \propto \delta R$.

1. Reducing invariants

Let us begin by invariants of form F^2 . The first three are (21):

$$\begin{aligned} I_1^F &= R_{\sigma\rho\chi\kappa} R^{\sigma\rho\chi\kappa} \\ I_2^F &= R_{\sigma\rho\chi\kappa} R^{\sigma\chi\rho\kappa} \\ I_3^F &= R_{\sigma\rho\chi\kappa} R^{\chi\kappa\sigma\rho} . \end{aligned}$$

As a consequence of the first Bianchi identity (26) and the skewsymmetries, the curvature tensor obey:

$$R_{\sigma\rho\chi\kappa} = R_{\chi\kappa\sigma\rho} . \quad (27)$$

Then,

$$I_3^F = R_{\sigma\rho\chi\kappa} R^{\chi\kappa\sigma\rho} = R_{\sigma\rho\chi\kappa} R^{\sigma\rho\chi\kappa} = I_1^F ,$$

while for I_2^F one finds,

$$I_2^F = R_{\sigma\rho\chi\kappa} R^{\sigma\chi\rho\kappa} = -(R_{\rho\chi\sigma\kappa} + R_{\chi\sigma\rho\kappa}) R^{\sigma\chi\rho\kappa} = -R_{\chi\rho\sigma\kappa} R^{\chi\sigma\rho\kappa} + R_{\chi\sigma\rho\kappa} R^{\chi\sigma\rho\kappa} = -I_2^F + I_1^F ,$$

$$2I_2^F = I_1^F ,$$

which let us with only one invariant of this kind, I_1^F .

Now, we translate the trace-like invariants (22) in a geometrical form:

$$\begin{aligned} I_1^{TrF} &= R_{\rho\mu\nu}{}^\rho R_\sigma^{\mu\nu\sigma} \\ I_2^{TrF} &= R_{\rho\mu\nu}{}^\rho R_\sigma^{\nu\mu\sigma} . \end{aligned}$$

Since the Ricci tensor $R_{\mu\nu} \equiv R_{\rho\mu\nu}{}^\rho$ is symmetric,² we have in fact only one invariant, $I_1^{TrF} = R_{\mu\nu} R^{\mu\nu}$.

At last, the only invariant of double traced form in F is:

$$I^{TrTrF} = R .$$

² Which is a consequence of the first Bianchi identity.

Analogously, in view of the Bianchi identities, only four invariants of the type G^2 remains (see appendix A):

$I_1^G = \delta_\beta R_{\sigma\rho\chi\kappa} \delta^\beta R^{\sigma\rho\chi\kappa}$	$TrTr\mathcal{G}_2 = \delta^\rho R_{\rho\chi} \delta_\mu R^{\mu\chi}$
$Tr\mathcal{G}_{10} = \delta_\beta R_{\sigma\chi} \delta^\chi R^{\sigma\beta}$	$Tr\mathcal{G}_{14} = \delta_\beta R_{\sigma\chi} \delta^\beta R^{\sigma\chi}$

V. GAUGE INVARIANCE CONDITION

With the invariants constructed above we collect seven types of Lagrangians for the gravitational field:

Lagr.	Inv.	Gauge Form	Geom. Form
$L_0^{(R_1)}$	$(I^{TrTrF})^n$	$(F_{ba}^{ab})^n$	R^n , $n = 1, 2$
$L_0^{(R_2)}$	I_1^{TrF}	$F_{b\mu a}^a F_c^{b\mu c}$	$R_{\mu\nu} R^{\mu\nu}$
$L_0^{(R_3)}$	I_1^F	$F_{\mu\nu}^{ab} F_{ab}^{\mu\nu}$	$R_{\alpha\beta\rho\sigma} R^{\alpha\beta\rho\sigma}$
$L_0^{(G_1)}$	$TrTr\mathcal{G}_2$	$G_{ab\beta}^{\beta a} G_{cb\mu}^{cb\mu}$	$\delta^\rho R_{\rho\chi} \delta_\mu R^{\mu\chi}$
$L_0^{(G_2)}$	$Tr\mathcal{G}_{14}$	$G_{\mu a\sigma}^{ab} G_{cb}^{\mu c\sigma}$	$\delta_\beta R_{\sigma\chi} \delta^\beta R^{\sigma\chi}$
$L_0^{(G_3)}$	$Tr\mathcal{G}_{10}$	$G_{a\sigma}^{abe} G_{deb}^{d\sigma}$	$\delta_\beta R_{\sigma\chi} \delta^\chi R^{\sigma\beta}$
$L_0^{(G_4)}$	I_1^G	$G_{\mu\nu\lambda}^{ab} G_{ab}^{\mu\nu\lambda}$	$\delta_\beta R_{\sigma\rho\chi\kappa} \delta^\beta R^{\sigma\rho\chi\kappa}$

(28)

We are considering Lagrangians only up to quadratic order in F and or G , which also include the linear invariant $I^{TrTrF} = R$ and their square R^2 . Actually, one can observe that if any invariant fulfills the gauge invariance condition, then any of its power will also do it, since this condition is linear in the derivatives $\frac{\partial L_0}{\partial F}$ and $\frac{\partial L_0}{\partial G}$. For instance,

$$L_0 = I^n, \quad \frac{\partial L_0}{\partial F} = n I^{n-1} \frac{\partial I}{\partial F}.$$

Therefore,

$$\frac{\partial I}{\partial F} [\dots] F = 0 \Rightarrow \frac{\partial L_0}{\partial F} [\dots] F = 0,$$

and the same follows for G .

Using the skewsymmetry $\nu \leftrightarrow \gamma$ of the equation (19) and the symmetry properties of the Riemann tensor, one can easily verify that all Lagrangians densities listed in (28) accomplish the gauge invariance condition. Then, any function of these invariants expressible in a Taylor series also will fulfill the gauge invariance condition.

VI. EQUATIONS OF MOTION

Here we will concentrate our attention on the effect of the term

$$L_0^{(G_1)} = \frac{1}{2} h h^a{}_\sigma h_c{}^\nu G_{ab}{}^\beta{}_\sigma G^{cb\mu}{}_{\mu\nu} = \frac{1}{8} h \delta^\rho R \delta_\rho R$$

on a gravitational theory based on the Einstein-Hilbert action plus the $L_0^{(G_1)}$ term. This Lagrangian density is equivalent, by the Bianchi identity, to the form $\frac{1}{2} h \delta^\rho R_{\rho\chi} \delta_\mu R^{\mu\chi}$, which is clearly analogous to Podolsky's second order term for Electrodynamics ($L_{Podolsky} \propto \partial^\rho F_{\rho\chi} \partial_\mu F^{\mu\chi}$). The choice of the particular Lagrangian $L_0^{(G_1)}$ is mainly motivated by this analogy. Besides this, the $L_0^{(G_1)}$ term also can be viewed as a kind of kinetic term for the scalar curvature, what approximate such description to the usual scalar fields. Moreover, this scalar is, up a surface term, present in the Schwinger-DeWitt renormalized effective action for an scalar field on a curved background [16]. Therefore, the field theory constructed over basis on $L_0^{(G_1)}$ can be considered an effective gravitational theory.

Taking a functional variation of the tetrad field, one finds:

$$h = \sqrt{-g} \ , \quad \delta h = \frac{1}{2} h g^{\lambda\nu} \delta g_{\lambda\nu} = h g^{\lambda\nu} h^a{}_\lambda \eta_{ab} \delta h^b{}_\nu$$

and

$$\begin{aligned} \delta L_0^{(G_1)} &= \frac{1}{4} \partial_\rho (h \partial^\rho R \delta R) - \frac{1}{4} \delta R \partial_\rho (h \partial^\rho R) + \\ &+ \frac{1}{4} h \left[\frac{1}{2} g^{\lambda\nu} \partial^\rho R \partial_\rho R - g^{\mu\nu} g^{\rho\lambda} \partial_\mu R \partial_\rho R \right] h^a{}_\lambda \eta_{ab} \delta h^b{}_\nu . \end{aligned}$$

On calculating the equations of motion, we must give special attention to the last term involving

$$\begin{aligned} \delta R &= -2 R_{\mu\beta} g^{\mu\nu} g^{\beta\lambda} h^a{}_\lambda \eta_{ab} \delta h^b{}_\nu + \\ &+ \frac{1}{h} \partial_\alpha \left[h \left(g^{\mu\nu} \delta \Gamma_{\mu\nu}^\alpha - g^{\nu\alpha} \delta \Gamma_{\nu\beta}^\beta \right) \right] , \end{aligned}$$

which will include several integration by parts. After these integrations and some cumbersome calculations, one finds:

$$\delta L_0^{(G_1)} = \frac{1}{2} \partial_\theta \mathcal{V}^\theta + \frac{1}{2} h \left[\delta_\lambda \delta_\nu [\diamond R] + \frac{1}{2} \delta_\lambda R \delta_\nu R - R_{\lambda\nu} \diamond R - g_{\lambda\nu} \diamond [\diamond R] - \frac{1}{4} g_{\lambda\nu} \delta^\rho R \delta_\rho R \right] h_a{}^\lambda \eta^{ab} \delta h_b{}^\nu ,$$

where

$$\begin{aligned} \mathcal{V}^\theta \equiv & -\frac{1}{2} \left(g^{\mu\nu} \delta \Gamma_{\mu\nu}^\theta - g^{\nu\theta} \delta \Gamma_{\nu\beta}^\beta \right) \partial_\rho (h \partial^\rho R) - \frac{1}{2} h \partial^\theta R \delta R + \\ & + \frac{1}{4} h \left(g^{\mu\nu} g^{\alpha\beta} - g^{\nu\alpha} g^{\mu\beta} \right) \delta_\alpha [\diamond R] \left(\delta^\theta_\nu \delta^\lambda_\mu \delta^\eta_\beta + \delta^\theta_\mu \delta^\lambda_\beta \delta^\eta_\nu - \delta^\theta_\beta \delta^\lambda_\nu \delta^\eta_\mu \right) \delta g_{\lambda\eta} , \end{aligned}$$

and

$$\diamond \equiv \delta_\beta \delta^\beta$$

is the Laplace-Beltrami operator on the Riemannian space.

Therefore, the second order contribution to the equation of motion will be

$$H_\nu^b \equiv h^{b\lambda} \delta_\lambda \delta_\nu [\diamond R] + \frac{1}{2} h^{b\lambda} \delta_\lambda R \delta_\nu R - R_\nu^b \diamond R - h^b_\nu \delta_\beta \delta^\beta [\diamond R] - \frac{1}{4} h^b_\nu \delta^\rho R \delta_\rho R . \quad (29)$$

Furthermore, if we include the usual first order Einstein-Hilbert and a matter Lagrangian densities,

$$S_T = \int d^n x \left(-\frac{hR}{2\chi} - \frac{\beta}{\chi} L_0^{(G_1)} + h \mathcal{L}_{matter} \right) ,$$

the field equations become

$$G^b_\nu + \beta H^b_\nu = \chi T^b_\nu , \quad (30)$$

or in a geometrical form,

$$R_{\lambda\nu} - \frac{1}{2} g_{\lambda\nu} R + \beta \left[\delta_\lambda \delta_\nu (\diamond R) + \frac{1}{2} \delta_\lambda R \delta_\nu R - R_{\lambda\nu} \diamond R - g_{\lambda\nu} \diamond (\diamond R) - \frac{1}{4} g_{\lambda\nu} \delta^\rho R \delta_\rho R \right] = \chi T_{\lambda\nu} ,$$

where G^b_ν is the Einstein tensor and

$$T_{\lambda\nu} \equiv \frac{2}{h} \frac{\delta (h \mathcal{L}_{matter})}{\delta g^{\lambda\nu}}$$

is the energy-momentum tensor of the matter fields written in terms of the metric field.

By analogy to the Alekseev-Arbuzov-Baikov [11], one could expect that the higher order terms, which can be until sixth derivative order, would be related to infrared corrections to General Relativity, giving sensible physical effects at large scales.

A. Covariant Conservation of $T_{\lambda\nu}$

Taking the covariant divergence of (30), we have

$$\delta^\nu G_{\nu\alpha} + \beta \delta^\nu H_{\nu\alpha} = \chi \delta^\nu T_{\nu\alpha} .$$

Now, from the first order case, we know that

$$\delta^\nu G_{\nu\alpha} \equiv 0 .$$

Applying the divergence to the equation (29), one finds

$$\begin{aligned} \delta^\nu H_{\nu\alpha} = & \delta^\nu \delta_\nu \delta_\alpha \Diamond R - g_{\nu\alpha} \delta^\nu \Diamond [\Diamond R] + \frac{1}{2} \delta_\nu R \delta^\nu \delta_\alpha R + \\ & + \frac{1}{2} \delta^\nu \delta_\nu R \delta_\alpha R - \delta^\nu R_{\nu\alpha} \Diamond R - R_{\nu\alpha} \delta^\nu \Diamond R - \frac{1}{4} g_{\nu\alpha} \delta^\nu (\delta^\rho R \delta_\rho R) . \end{aligned}$$

Using the commutation relation

$$[\delta_\nu, \delta_\alpha] A^\tau = R_{\alpha\nu}{}^{\tau\xi} A_\xi$$

and the second Bianchi identity, we arrive at

$$\delta^\nu H_{\nu\alpha} = R_{\alpha\xi} \delta^\xi \Diamond R - R_{\nu\alpha} \delta^\nu \Diamond R + \frac{1}{2} \delta_\nu R \delta^\nu \delta_\alpha R - \frac{1}{2} \delta^\rho R \delta_\alpha \delta_\rho R = 0 .$$

Then, the covariant conservation of $T_{\mu\nu}$ is established:

$$\delta^\mu (G_{\mu\nu} + \beta H_{\mu\nu}) \equiv 0 \implies \delta^\mu T_{\mu\nu} = 0 ,$$

as expected from the coordinate invariance of the Lagrangian density.

B. Static Isotropic Solution

In the case of an static isotropic metric,

$$ds^2 = e^{\nu(r)} dt^2 - e^{\lambda(r)} dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2$$

in the vacuum, the equations of motion (30) are reduced, in the linear approximation, to the following coupled linear equations:

$$\begin{aligned} & \nu'' + \frac{2}{r} \nu' + \\ & + \beta \left(\nu^{(6)} + \frac{6}{r} \nu^{(5)} - \frac{2}{r} \lambda^{(5)} - \frac{2}{r^2} \lambda^{(4)} + \frac{8}{r^3} \lambda''' - \frac{24}{r^4} \lambda'' + \frac{48}{r^5} \lambda' - \frac{48}{r^6} \lambda \right) = 0 , \\ & \frac{1}{2} \left(\frac{\lambda'}{r} - 2 \frac{\lambda}{r^2} + \frac{\nu'}{r} - \nu'' \right) + \\ & - \beta \left(\nu^{(6)} + \frac{3}{r} \nu^{(5)} - \frac{12}{r^2} \nu^{(4)} + 12 \frac{\nu'''}{r^3} - \frac{2}{r} \lambda^{(5)} + \frac{4}{r^2} \lambda^{(4)} + 8 \frac{\lambda'''}{r^3} - 48 \frac{\lambda''}{r^4} + 96 \frac{\lambda'}{r^5} - 96 \frac{\lambda}{r^6} \right) = 0 . \end{aligned}$$

To solve this system, we use the Frobenius method, based on a series expansion:

$$\nu(r) = \sum_n \nu_n r^{s+n}, \quad \lambda(r) = \sum_n \lambda_n r^{s+n}.$$

From the first terms in the series, we find $s = -1$, and the recursion relations above:

$$\begin{aligned} \lambda_{n+4} &= \frac{\nu_n (n-2)}{4\beta (n+4) (n+2) (n+1) (n-\frac{1}{2})}, \\ \nu_{n+4} &= -\frac{\nu_n}{2\beta (n+4) (n+3) (n+2) (n-\frac{1}{2})}, \end{aligned}$$

such that the solution can be written as

$$\begin{aligned} \nu(r) &= \sum_{m=0}^3 \nu_m r^{m-1} \left(1 + \sum_{n=0}^{\infty} c_{nm} \right), \\ \lambda(r) &= \sum_{m=0}^3 \lambda_m r^{m-1} - \sum_{m=0}^3 \nu_m r^{m-1} \sum_{n=0}^{\infty} \frac{(4n+m-2)(4n+3+m)}{2(4n+1+m)} c_{nm}, \end{aligned}$$

where ν_m and λ_m , with $m \in \{0, 1, 2, 3\}$, are the integration constants specified by the boundary conditions, and

$$c_{nm} \equiv \left(-\frac{r^4}{2\beta} \right)^{n+1} \frac{(m+1)!}{(4n+m+4)!} \frac{(4n+m+1)!!!!}{(m+1)!!!!} \frac{(m-\frac{9}{2})!!!!}{(4n+m-\frac{1}{2})!!!!}$$

The notation $a!!!!$ stands for:

$$(a+4)!!!! = (a+4) . a!!!! .$$

The convergence of the series, tested by the ratio test,

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{\nu_{n+1}}{\nu_n} \right| &= \left| \frac{r^4}{2\beta} \right| \lim_{n \rightarrow \infty} \left| D_{n,m} \frac{1}{(4n+m+7)} \right| = 0, \\ \lim_{n \rightarrow \infty} \left| \frac{\lambda_{n+1}}{\lambda_n} \right| &= \left| \frac{r^4}{2\beta} \right| \lim_{n \rightarrow \infty} \left| D_{nm} \frac{(4n+m+2)(4n+m+1)}{(4n+m+5)(4n+m+3)(4n+m-2)} \right| = 0, \\ D_{nm} &\equiv \frac{1}{(4n+m+8)(4n+m+6)(4n+m+\frac{7}{2})}, \end{aligned}$$

shows that both are convergent with an infinite radius of convergence.

Therefore, in the first order approximation for β , we have

$$\begin{aligned} \nu(r) &= \frac{\nu_0}{r} \left(1 + \frac{1}{24\beta} r^4 \right) + \nu_1 \left(1 - \frac{1}{60\beta} r^4 \right) + \\ &\quad + \nu_2 r \left(1 - \frac{1}{360\beta} r^4 \right) + \nu_3 r^2 \left(1 - \frac{1}{1050\beta} r^4 \right) + \mathcal{O}(\beta^2) \\ \lambda(r) &= -\frac{\nu_0}{r} + \lambda_1 + \lambda_2 r + \lambda_3 r^2 + \frac{1}{6\beta} \left(-\frac{\nu_0}{4} r^3 + \frac{\nu_1}{10} r^4 + \frac{\nu_2}{60} r^5 + \frac{\nu_3}{175} r^6 \right) + \mathcal{O}(\beta^2) \end{aligned} \tag{31}$$

An analysis of the solution (31) reveals the expected weak field behaviour at short scales ($\frac{\nu_0}{r}$), and deviation from this for a mesoscale, since we are dealing only with the linear approximation. Correspondingly, we find $\nu_0 = 2GM/c^2$ where M is the mass of the central body.

The remaining integration constants set scale distances where modifications of the Newtonian behaviour appear. For instance, consider the Einstein-Hilbert theory with cosmological constant. The static spherically symmetric solution is

$$\nu(r) = -\lambda(r) = 1 - \frac{2GM}{c^2} \frac{1}{r} - \frac{\Lambda}{3} r^2$$

where the cosmological constant sets an scale distance given by the de Sitter pseudo-radius.

Analogously, in our case, the ν_1 constant sets a constant potential, which can be a mean nonlocal value of the effective Lagrangian proposed, ν_2 sets an scale distance where a constant mean force appears, and ν_3 represents a gradient of force, in the same way as the cosmological constant in the example above. Similar reasoning can be developed for the other constants in the model.

The contribution of each constant to the net force could be fixed by requiring that it fits the observational data for the tests of the gravitation. This task deserves a careful investigation of its own and is presently under investigation by the authors by means of the study of galaxy rotation curves, geodesic motion, perihelion shift, gravitational lenses and redshift.

VII. CONCLUSION

We have applied the second order gauge theory [10] to the local gauge theory for the homogeneous Poincaré group. It was found that the geometrical counterparts of the usual field strength F and the second order field strength $G = DF$ are the Riemann tensor R and its (space-time) covariant derivative δR . It followed the analysis of the second order invariants composed with the geometrical entities.

We demonstrate – employing the symmetry properties of the curvature tensor – that the only independent Lagrangian densities for the gravitational field in a Riemannian manifold of arbitrary dimension are the seven ones listed in table (28). Linear combinations of terms proportional to powers of R , as the familiar quadratic term in the curvature, are of first

order in the gauge potential ω , therefore, in the context of the second order gauge theory, the contributions of second order in the Lagrangian density, which are those including second derivatives of the gauge potential, are of type δR .

We derived equations of motion using a particularly simple choice for the second order gauge Lagrangian inspired in the Podolsky's proposal for a Generalized Electrodynamics. We found the static isotropic solution of these equations in the linear approximation, showing that at short distances the gravitational field behaves exactly as the Newton's law, but at meso-large distance scales the higher order contribution dominates, exhibiting a modified potential.

In the future, we will study other solutions of these field equations, searching for massive modes which do not violate the local gauge symmetry. Our guide in these calculations shall be the treatment given in [10] to the $U(1)$ case, where an effective mass for the photon was derived. To do this, one naturally must concern about the determination of the conserved current associated with the local Lorentz symmetry and the relationship to the global diffeomorphic invariance of the theory.

Another perspective is to apply the second order equations of motion (30) to a Friedmann-Robertson-Walker metric. The goal is to seek for a accelerated regimes of the cosmological model arising from the higher order terms. This proposal is now under investigation.

ACKNOWLEDGEMENTS

R. R. C. and L. G. M. are grateful to Fundação de Amparo à Pesquisa do Estado de S. Paulo (FAPESP) for support (grants 02/05763-8 and 02/10263-4 respectively); P. J. P. thanks to CTA staff. C. A. M. M. thanks the hospitality of the International Centre of Theoretical Physics, under Federation Arrangement, where part of this work was done. The authors acknowledge Prof. B. M. Pimentel, Prof. R. Aldrovandi and Prof. A. J. Accioly for the continuing incentive to this project, and an unknown referee for useful comments.

APPENDIX A: APPENDIX: COUNTING SECOND ORDER INVARIANTS

1. Counting GG Invariants

First, let us analyse how many are the possible contractions of kind GG . This is done by means of tables as in the section IV A. The first one is constructed fixing, for instance, the last index:

Fix. e	Fix. a	Fix. b	Fix. c	Fix. d
cyclic	$(abcd)$	$(bacd)$	$(cabd)$	$(dabc)$
	$(acdb)$	$(bcda)$	$(cbda)$	$(dbca)$
	$(adbc)$	$(bdac)$	$(cdab)$	$(dcab)$
non-cycl.	$(abdc)$	$(badc)$	$(cadb)$	$(dacb)$
	$(acbd)$	$(bcad)$	$(cbad)$	$(dbac)$
	$(adcb)$	$(bdca)$	$(cdba)$	$(dcba)$

Analogous tables result when we fix the indices d, c, b and a . For each table, non-cyclic permutations are equivalent to cyclic ones, giving:

Fix. e	Fix. a	Fix. b	Fix. c	Fix. d
cyclic	$(abcd)$	—	—	—
	$(acdb)$	$(bcda)$	—	—
	$(adbc)$	$(bdac)$	$(cdab)$	—

an similarly for the other four tables.

Using the cyclic permutation symmetry, one can identify elements of different tables, reducing the number of invariants. By the skew-symmetry in the first G and renaming dummy indices, it follows:

$\mathcal{G}_1 = [abcde] [abcde]$	$\mathcal{G}_6 = [abcde] [cdbea]$
$\mathcal{G}_2 = [abcde] [beacd]$	$\mathcal{G}_7 = [abcde] [adbec]$
$\mathcal{G}_3 = [abcde] [adceb]$	$\mathcal{G}_8 = [abcde] [acbde]$
$\mathcal{G}_4 = [abcde] [aecbd]$	$\mathcal{G}_9 = [abcde] [acdeb]$
$\mathcal{G}_5 = [abcde] [debac]$	$\mathcal{G}_{10} = [abcde] [abdce]$

One can further apply the cyclic permutation symmetry to the first G in these remaining

invariants and reduce even more the number of independent quantities. Beginning with $\mathcal{G}_{10}$:

$$\mathcal{G}_{10} = -([abdec] + [abecd])[abdce] = \mathcal{G}_1 - \mathcal{G}_{10} \Rightarrow 2\mathcal{G}_{10} = \mathcal{G}_1 .$$

On the other hand, for $\mathcal{G}_9$:

$$\mathcal{G}_9 = -([abdec] + [abecd])[acdeb] = 2\mathcal{G}_4 + \mathcal{G}_9 \Rightarrow \mathcal{G}_4 = 0 .$$

Proceeding in the same way, one finds the following identities:

$$\begin{aligned} 2\mathcal{G}_{10} &= \mathcal{G}_1; & 2\mathcal{G}_6 &= \mathcal{G}_5 ; \\ \mathcal{G}_2 &= \mathcal{G}_3 = \mathcal{G}_4 = \mathcal{G}_7 = \mathcal{G}_8 = \mathcal{G}_9 = 0 . \end{aligned}$$

which give two independent invariants,

$$I_1^G = [abcde][abcde] , \quad I_2^G = [abcde][debac] .$$

2. Counting $(TrG)^2$ Invariants

Starting with the three independent traces listed in (23), and considering skew-symmetries, the possible quadratic combinations are:

$Tr\mathcal{G}_1 = [\cdot a \cdot bc][\cdot a \cdot bc]$	$Tr\mathcal{G}_{10} = [\cdot ab \cdot c][\cdot ba \cdot c]$
$Tr\mathcal{G}_2 = [\cdot a \cdot bc][\cdot b \cdot ac]$	$Tr\mathcal{G}_{11} = [\cdot ab \cdot c][\cdot bc \cdot a]$
$Tr\mathcal{G}_3 = [\cdot ab \cdot c][\cdot a \cdot bc]$	$Tr\mathcal{G}_{12} = [\cdot ab \cdot c][\cdot ca \cdot b]$
$Tr\mathcal{G}_4 = [\cdot ab \cdot c][\cdot b \cdot ac]$	$Tr\mathcal{G}_{13} = [\cdot ab \cdot c][\cdot cb \cdot a]$
$Tr\mathcal{G}_5 = [\cdot ab \cdot c][\cdot c \cdot ab]$	$Tr\mathcal{G}_{14} = [\cdot ab \cdot c][ab \cdot \cdot c]$
$Tr\mathcal{G}_6 = [ab \cdot \cdot c][\cdot a \cdot bc]$	$Tr\mathcal{G}_{15} = [\cdot ab \cdot c][ac \cdot \cdot b]$
$Tr\mathcal{G}_7 = [ab \cdot \cdot c][\cdot c \cdot ab]$	$Tr\mathcal{G}_{16} = [\cdot ab \cdot c][bc \cdot \cdot a]$
$Tr\mathcal{G}_8 = [\cdot ab \cdot c][\cdot ab \cdot c]$	$Tr\mathcal{G}_{17} = [ab \cdot \cdot c][ab \cdot \cdot c]$
$Tr\mathcal{G}_9 = [\cdot ab \cdot c][\cdot ac \cdot b]$	$Tr\mathcal{G}_{18} = [ab \cdot \cdot c][ac \cdot \cdot b]$

The last two invariants can not be converted into any other using the symmetries at our disposal. Each one of the preceding $Tr\mathcal{G}$ must be analyzed, case by case, in a search for eventual interdependence.

Take, for example, the 16th term, and rewrite it as bellow:

$$Tr\mathcal{G}_{16} = -[\cdot acb \cdot][bc \cdot \cdot a] - [\cdot a \cdot cb][bc \cdot \cdot a] \Rightarrow 2Tr\mathcal{G}_{16} = Tr\mathcal{G}_7 .$$

Repeat the reasoning for, say, the 15th invariant:

$$Tr\mathcal{G}_{15} = -[\cdot acb\cdot][ac\cdot b] - [a\cdot cb][ac\cdot b] = Tr\mathcal{G}_{14} - Tr\mathcal{G}_6 .$$

As soon as we perform this same check for all the above invariants, only eight of them are kept:

$Tr\mathcal{G}_3 = [\cdot ab\cdot c][\cdot a\cdot bc]$	$Tr\mathcal{G}_{11} = [\cdot ab\cdot c][\cdot bc\cdot a]$	(A1)
$Tr\mathcal{G}_5 = [\cdot ab\cdot c][\cdot c\cdot ab]$	$Tr\mathcal{G}_{14} = [\cdot ab\cdot c][\cdot ab\cdot c]$	
$Tr\mathcal{G}_6 = [ab\cdot\cdot c][\cdot a\cdot bc]$	$Tr\mathcal{G}_{17} = [ab\cdot\cdot c][ab\cdot\cdot c]$	
$Tr\mathcal{G}_{10} = [\cdot ab\cdot c][\cdot ba\cdot c]$	$Tr\mathcal{G}_{18} = [ab\cdot\cdot c][ac\cdot\cdot b]$	

3. Counting $(TrTrG)^2$ Invariants

From $T_{abc}^{(1)} \equiv [\cdot a\cdot bc]$ one can take a trace again:

$$T_c^{(1)} \equiv [\cdot \circ \circ c] .$$

From $T_{abc}^{(2)} \equiv [\cdot ab\cdot c]$ one finds $T_c \equiv [\cdot \circ \circ \cdot c]$ which can be reduced to $T_c^{(1)}$ using the G skewsymmetry in the first two indexes and changing dummy indexes. Another possible trace is constructed from $T_{abc}^{(2)}$:

$$T_b^{(2)} \equiv [\cdot \circ b \cdot \circ] . \quad (A2)$$

But it also is not independent of $T_c^{(1)}$:

$$T_b^{(2)} \equiv [\cdot \circ b \cdot \circ] = -[\cdot \circ \circ b] - [\cdot \circ \circ b] = -T_b^{(1)} - [\circ \circ \circ b] = -2T_b^{(1)} .$$

Let us set $T_b^{(2)}$ as the independent double trace.

There are an internal double trace of $T_{abc}^{(3)} \equiv [ab\cdot\cdot c]$ which is independent of $T_c^{(2)}$:

$$T_b^{(3)} \equiv [\circ b \cdot \cdot \circ] . \quad (A3)$$

The other double trace of $T_{abc}^{(3)}$ is,

$$T_b \equiv [b \circ \cdot \cdot \circ] = -T_b^{(3)} .$$

Then, we have the following set of independent double traces:

$$\begin{aligned} TrTr\mathcal{G}_1 &= [\cdot \circ b \cdot \circ][\cdot \circ b \cdot \circ] \\ TrTr\mathcal{G}_2 &= [\circ b \cdot \cdot \circ][\circ b \cdot \cdot \circ] \\ TrTr\mathcal{G}_3 &= [\cdot \circ b \cdot \circ][\circ b \cdot \cdot \circ] . \end{aligned} \quad (A4)$$

4. Reducing the G^2 invariants using Bianchi identities

Consider the reduction of the number of quadratic invariants in G by means of Bianchi identities. Using the geometric form, the first two invariants are:

$$I_1^G = \delta_\beta R_{\sigma\rho\chi\kappa} \delta^\beta R^{\sigma\rho\chi\kappa} , \quad I_2^G = \delta_\beta R_{\sigma\rho\chi\kappa} \delta^\chi R^{\beta\kappa\sigma\rho} .$$

Applying the second Bianchi identity to I_2^G we have:

$$\begin{aligned} I_2^G &= -\delta_\beta R_{\sigma\rho\chi\kappa} (\delta^\chi R^{\kappa\beta\rho\sigma} + \delta^\beta R^{\chi\kappa\rho\sigma}) = I_1^G - I_2^G \Rightarrow \\ &\Rightarrow 2I_2^G = I_1^G , \end{aligned}$$

therefore it is sufficient to consider only I_1^G .

Let us analyse now the trace invariants in G , (24):

$Tr\mathcal{G}_3 = \delta_\beta R_{\sigma\zeta\chi}{}^\zeta \delta_\mu R^{\sigma\beta\chi\mu}$	$Tr\mathcal{G}_{11} = \delta_\beta R_{\sigma\chi} \delta^\sigma R^{\chi\beta}$
$Tr\mathcal{G}_5 = \delta_\beta R_{\sigma\rho\chi}{}^\rho \delta_\mu R^{\beta\chi\sigma\mu}$	$Tr\mathcal{G}_{14} = \delta_\beta R_{\sigma\chi} \delta^\beta R^{\sigma\chi}$
$Tr\mathcal{G}_6 = \delta^\rho R_{\sigma\rho\chi\zeta} \delta_\kappa R^{\sigma\chi\zeta\kappa}$	$Tr\mathcal{G}_{17} = \delta^\rho R_{\sigma\rho\chi\kappa} \delta_\mu R^{\sigma\mu\chi\kappa}$
$Tr\mathcal{G}_{10} = \delta_\beta R_{\sigma\chi} \delta^\chi R^{\sigma\beta}$	$Tr\mathcal{G}_{18} = \delta^\rho R_{\sigma\rho\chi\kappa} \delta_\mu R^{\chi\mu\sigma\kappa}$

Comparing $Tr\mathcal{G}_{10}$ with $Tr\mathcal{G}_{11}$ one sees that both are the same invariant, due to the symmetry of Ricci tensor.

Using the second Bianchi identity, it follows:

$$Tr\mathcal{G}_3 = \delta_\beta R_{\sigma\chi} g_{\mu\rho} (\delta^\beta R^{\rho\sigma\chi\mu} + \delta^\sigma R^{\beta\rho\chi\mu}) = Tr\mathcal{G}_{14} - Tr\mathcal{G}_{10}$$

and in the same way,

$$Tr\mathcal{G}_5 = Tr\mathcal{G}_6 = -\frac{1}{2}Tr\mathcal{G}_{17} = -Tr\mathcal{G}_{18} = -Tr\mathcal{G}_3 .$$

This shows that only $Tr\mathcal{G}_{14}$ and $Tr\mathcal{G}_{10}$ can be hold independent.

We apply the same technique to the double traced invariants (25):

$$\begin{aligned} TrTr\mathcal{G}_1 &= \delta_\beta R \delta^\beta R \\ TrTr\mathcal{G}_2 &= \delta^\rho R_{\rho\chi} \delta_\mu R^{\mu\chi} \\ TrTr\mathcal{G}_3 &= -\delta_\beta R \delta_\mu R^{\mu\beta} . \end{aligned}$$

The second Bianchi identity shows us that we have only one invariant in such case:

$$TrTr\mathcal{G}_3 = \delta^\rho_\zeta g^{\sigma\chi} \left(\delta_\sigma R_{\beta\rho\chi}{}^\zeta + \delta_\rho R_{\sigma\beta\chi}{}^\zeta \right) \delta_\mu R^{\mu\beta} = -2TrTr\mathcal{G}_2 = -\frac{1}{2}TrTr\mathcal{G}_1 .$$

- [1] P. Astier *et al.*, Astron. Astrophys. **447**, 31 (2006); A. G. Riess *et al.*, Astrophys. J. **607**, 665 (2004).
- [2] S. Nojiri and S. D. Odintsov, *Modified $f(R)$ gravity consistent with realistic cosmology: from matter dominated epoch to dark energy universe*, available as [hep-th/0608008]; S. Capozziello, Int. J. Mod. Phys. **D11**, 483 (2002); S. M. Carroll, V. Duvvuri, M. Trodden and S. Turner, Phys. Rev. **D70**, 043528 (2004); U. Gunther, A. Zhuk, V. Bezerra and C. Romero, Class. Quant. Grav. **22**, 313 (2005).
- [3] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press (1984); R. Casana, C. A. M. de Melo and B. M. Pimentel, Proc.Sci. **WC2004** (2004) 013; R. Casana, C. A. M. de Melo and B. M. Pimentel, *Braz. J. Phys.* **35**, 1151 (2005); R. Casana, C. A. M. de Melo and B. M. Pimentel, *Spinorial Field and Lyra Geometry*, Astrophys. Sp. Sc. (2006), DOI: 10.1007/s10509-006-9048-5; available online.
- [4] B. Podolsky, Phys. Rev. **62**, 68 (1942); B. Podolsky and C. Kikuchi, Phys. Rev. **65**, 228 (1944); B. Podolsky and P. Schwed, Rev. Mod. Phys. **20**, 40 (1948).
- [5] K. S. Stelle, Gen. Rel. Grav. **9**, 353 (1978); A. Accioly, A. Azeredo and H. Mukai, J. Math. Phys. **43**, 473 (2002).
- [6] R. P. Feynman, *Feynman Lectures on Gravitation*, Westview Press (2002).
- [7] M. H. Goroff and A. Sagnotti, Phys. Lett. **160B**, 81 (1985).
- [8] B. S. DeWitt, *Dynamical Theory of Groups and Fields*, Gordon & Breach Science Pub (1965); S. M. Christensen, Phys. Rev. **D14**, 2490 (1976).
- [9] R. Utiyama, Phys. Rev. **101**, 1597 (1956).
- [10] R. R. Cuzinatto, C. A. M. de Melo and P. J. Pompéia, *Second order gauge theory*, Ann. Phys. **322**, 1211 (2007).
- [11] A. I. Alekseev, B. A. Arbuzov and V. A. Baikov, Theor. Math. Phys. **52**, 739 (1982).
- [12] S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummings, *Normal forms for tensor polynomials: I. The Riemann tensor*, Class. Quantum Grav. **9**, 1151 (1992).

- [13] Y. Décanini and A. Folacci, *Standard forms for the metric variations of the action terms of sixth-order gravity and approximated stress-energy tensor*, available as arXiv: 0706.0691 (2007).
- [14] M. Blagojevic, *Gravitation & Gauge Symmetries*, Taylor & Francis (2001).
- [15] V. De Sabbata and M. Gasperini, *Introduction to Gravitation*, World Scientific (1985).
- [16] I. G. Avramidi, *Heat Kernel and Quantum Gravity*, Berlin, Springer (2000).